

RESEARCH

Open Access



Differences between problematic internet and smartphone use and their psychological risk factors in boys and girls: a network analysis

Dmitri Rozgonjuk^{1,2}, Lukas Blinka^{3*}, Nana Löchner^{1*}, Anna Faltýnková³, Daniela Husarova⁴ and Christian Montag¹

Abstract

Background Problematic internet and smartphone use are significant health challenges for contemporary adolescents. However, their mutual relationship is unclear because studies investigating these phenomena are scarce. The present study aimed to investigate the psychological risks and protective factors associated with problematic internet and smartphone use.

Method A representative sample of Slovak adolescents ($N = 4070$, $M_{age} = 14.38$, $SD_{age} = 0.77$, 50.5% girls) from the Health Behavior in School-aged Children project was analyzed using network analysis separately for boys and girls.

Results The results showed weak (for boys) and moderate (for girls) associations between problematic internet use and problematic smartphone use. Risk factors showed stronger associations with problematic internet use than problematic smartphone use, with the exception of fear of missing out, which was strongly associated with problematic smartphone use. The central nodes were externalizing problems for boys and internalizing problems, externalizing problems, and resilience for girls.

Conclusion The study concluded that while problematic internet use and problematic smartphone use are somewhat related, they differ at the psychological level. In addition, the phenomena are rather different between boys and girls.

Keywords Problematic internet use, Problematic smartphone use, Psychology risk factors, HBSC, Network analysis

Introduction

Over the past two decades, increasing access to digital technology has transformed the lives of young people worldwide. Modern adolescents work with the internet and other digital technologies on an intensive daily basis [1]. The expansion and constant accessibility of internet and other technologies have created great opportunities for learning, work, entertainment, and personal exploration and growth [2]. However, intense online technology use may lead to various social and health risks, including reduced sleep quality [3], obesity [4], and reduced academic performance [5]. Concerns regarding excessive and potentially addictive use have been repeatedly

*Correspondence:

Lukas Blinka
lukasblinka@gmail.com

Nana Löchner
nana.loechner@uni-ulm.de

¹ Department of Molecular Psychology, Institute of Psychology and Education, Ulm University, Helmholtzstraße 8/1, 89081 Ulm, Germany

² Institute of Mathematics and Statistics, University of Tartu, Tartu, Estonia

³ Psychology Research Institute, Faculty of Social Studies, Masaryk University, Joštova 10, 60200 Brno, Czech Republic

⁴ Department of Health Psychology and Methodology Research, Faculty of Medicine, P.J. Safarik University in Kosice, Košice, Slovakia



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

expressed [6, 7]. Of the many different forms of excessive technology use, considerable attention has been paid to problematic internet use (PIU) and problematic smartphone use (PSU). Much research has already been devoted to these two phenomena in adolescence [8, 9], but the relationship between them (especially with regard to adolescents) has been studied to a limited extent. Our study fills this gap by comparing the similarities and differences between PIU and PSU in terms of their risk and protective factors (i.e., psychological variables that might affect susceptibility to PIU or PSU). Our focus is specifically on adolescent users, since they are often seen as a particularly vulnerable group in terms of the development of problematic forms of internet and smartphone use [1, 10].

Both PIU and PSU are operationalized as the inability to control one's use of the medium, which leads to harmful consequences and disruptions in daily functioning [11, 12]. Even when facing these negative consequences, users have a diminished capacity to limit their time spent in the medium, and they are preoccupied with it, even when not online [13]. The term PIU covers a large number of excessive online activities, especially online gaming, social networking sites use (SNS), chatting, video watching, or online shopping [12]. Smartphones are internet-enabled devices providing instant and nearly unlimited access to online activities. In principle, when using a smartphone, the user is almost always simultaneously connected to the internet. Thus, in terms of user patterns, it can be assumed that the two phenomena overlap to a certain extent. On the other hand, the application use may be slightly different, because some social media use (e.g., Instagram, WhatsApp) is optimized for smartphones. The existing literature reports weak to strong correlations between PSU and PIU (e.g., $r=0.21$ in Choi et al. [14]; $r=0.40$ in Kwon et al. [15]; $r=0.50$ in Lachmann et al. [16]; $r=0.64$ in Škařupová et al. [17]). Although positive associations have been demonstrated, several studies have pointed out differences in usage patterns, gender, personality traits, and psychological variables between these two types of problematic behavior. While PSU has been found to be especially related (and almost identical) to social media use [18], extreme PIU scores were found to be related to online gaming [19]. In other words, both phenomena rely on somewhat different need satisfaction and anticipated rewards [20]. Specifically, in terms of internet usage, boys were reported to be more prone to addictive use than girls, whereas this pattern was reversed for smartphone use [14, 21]. Furthermore, lower extraversion was associated with higher PIU but was unrelated to PSU, whereas lower openness to experience was linked to higher PSU but not to PIU [22]. These results suggest that while PIU and PSU are related,

a significant portion of unexplained variance remains that represents the differences between the constructs.

Both PIU and PSU are often studied in the context of users' psychological characteristics and susceptibility to developing problematic forms of use. Previous meta-analyses [23–25] have suggested that the most consistent risk factors stem from the following areas: (1) high impulsivity and attention/hyperactivity disorders; (2) negative emotionality, anxiety, and depressive symptoms; and (3) low self-esteem and self-directedness. On the other hand, resilience and high self-control are often reported as the most important protective factors in terms of PIU and PSU [26, 27].

Impulsivity is frequently linked to addictive behaviors [28]. Internet users with higher impulsivity present executive dysfunction and deficient inhibitory control, which may contribute to problems with online technology use [29]. Together with disrupted self-control, attention problems, aggression, hyperactivity, and impulsivity are part of the construct of externalizing problems [30]. According to previous research, externalizing problems are relatively common in adolescent problematic media users and they were found to be significantly associated with PIU, specifically excessive social media use or internet gaming addiction [31–34].

In addition to externalizing problems, adolescents may also develop internalizing problems that include affective states, such as anxiety, social withdrawal, depression [35], diminished self-esteem, and feelings of hopelessness—all of which have been identified as risk factors in terms of developing both PIU and PSU [36–39]. It has been suggested that individuals with negative emotionality may tend to use smartphones or the internet excessively as a coping mechanism to eliminate distress [40, 41].

Internalizing problems in adolescence often go hand-in-hand with social anxiety, withdrawal from peer relationships, lack of social competence, and shyness [42, 43]. Thus, these individuals experience problems with social functioning and social inclusion [44]. Adolescents who need to belong and whose social connections are unsatisfied in real life might tend to fulfill these needs through SNS [45]. Przybylski et al. proposed that fear of missing out (FoMO) may explain these dynamics. Fear of missing out is defined as a pervasive apprehension that others might have more rewarding experiences or acquire useful information that one does not have access to [46]. To not miss something important on the site, people with increased FoMO feel the need to be as often online as possible. They often experience anxiety offline and feel pressured to constantly check for new information [47]. Previous research has demonstrated robust associations between FoMO and problematic social networks use [48], PIU [49], and PSU [50]. A potential explanation to

these findings may lie in unmet social needs: it has been demonstrated that adolescents with higher FoMO also tend to have a higher need for popularity/belonging as well as higher social media use intensity [45].

Resilience is one of the strongest protective factors for PIU or PSU. Resilience is a multidimensional construct defined as the ability to adapt positively to life conditions and thrive, even in the face of adversity [51]. A resilient individual can use constructive coping strategies to successfully deal with adverse life events [52]. Resilience has been repeatedly suggested as a protective factor against various psychopathologies and risky behaviors, including internet, smartphone, and social media addiction [53–56]. Young people face many stressful challenges due to biological, psychological, and social changes. Adolescents with higher resilience have better internal resources to cope with stressful events, which might lead them to become less involved in using the internet to regulate negative emotions [57].

Although a relatively large body of work has examined PIU and PSU use separately, examinations of both are relatively scarce. Based on the literature review, we assume that they are related (e.g., share some of the predictors), but distinct phenomena. Owing to the different usage patterns of boys and girls, we further assumed that the differences would be reflected at the gender level as well. Thus, the aim of this study was to examine what psychological risk and protective factors are shared by PIU and PSU and what factors define the dividing line. With the help of a network analysis, we aim to examine the association separately for boys and girls.

Methods

Data collection and sample

In the present study, we used data from the WHO-collaborative Health Behavior in School-aged Children (HBSC) study [58]. The HBSC is a cross-sectional study carried out at 4-year intervals in 50 countries and regions across Europe and North America. Only the survey conducted in Slovakia in 2018 was analyzed because it used the key variables essential for the purpose of this study. A nationally representative sample of Slovak adolescents aged 11–15 was obtained using a two-step data collection procedure. First, the list of all eligible schools in Slovakia was obtained from the Slovak Institute of Information and Prognosis for Education, and then stratified by region and type of school (primary vs. secondary school). A total of 140 schools were randomly selected and asked to participate; and 109 agreed to participate (response rate 77.85%). In the second step, one class from each grade within the target age group was randomly selected from each school. Data were collected anonymously using self-report electronic questionnaires administered by trained

administrators during classroom sessions. Participation was voluntary, and passive parental informed consent was obtained before administering the questionnaires.

The sample included more than 8405 Slovakian adolescents aged 11–15. However, some of the key variables (including the PSU scale) were administered only to respondents aged 13–15, which reduced the sample size to 5053 participants. Moreover, we removed the data of students whose responses were missing for at least 75% of the scale items used in this study. Therefore, the effective sample, which comprised 4070 adolescents (age $M = 14.38$, $SD = 0.77$; 2013 boys and 2057 girls), was used in this study. Participants' gender was not associated with missing data. The average age of the excluded samples ($M = 14.24$) was lower than that of students who were not excluded from the dataset because of missing data ($M = 14.38$), $t(1384.6) = -4.915$, $p < 0.001$. Data for other study participants who had missing data were imputed (see the analysis section for details).

Measures

In the current study, we used data that included participants' sociodemographic characteristics and their responses to scales that assessed the severity of PIU and PSU. They also included experiencing externalizing and internalizing problems, fear of missing out, resilience, and hopelessness. Descriptive statistics for these scales and their internal consistencies are presented in Table 1.

PIU was measured using the Excessive Internet Use Scale [EIU; 59]. The scale consists of five items covering five of the six factors of the Griffiths component model of behavioral addiction [60]. These factors are salience (i.e., "I have gone without eating and sleeping because of the internet"), withdrawal symptoms (i.e., "I have felt bothered when I cannot be on the internet"), tolerance (i.e., "I have caught myself surfing when I am not really interested"), relapse (i.e., "I have tried unsuccessfully to spend less time on the internet"), and conflict (i.e., "I have spent less time than I should with either family, friends, or doing schoolwork because of the time I spend on the internet"). Participants used a 4-point scale (ranging from 1 = "never" to 4 = "very often") to express how often they had experienced certain symptoms in the preceding 12 months. The final variable was the sum of the five items.

PSU was measured using the Problematic Mobile Phone Use Scale [MPPUS-10; 61]. This is a shortened version of the Mobile Phone Problem Use Scale [62] and consists of 10 items (e.g., "I have used my mobile phone to make myself feel better when I was feeling down"). Participants answered on a 5-point scale (1 = "strongly disagree", 5 = "strongly agree") to what extent they agreed with each statement about their everyday mobile use. A

Table 1 Descriptive statistics of the total sample including boys and girls

Variable	Theoretical range	Total sample (N = 4070)			Boys (N = 2013)		Girls (N = 2057)		Boys–girls difference test		
		M	SD	α	M	SD	M	SD	W	p	d (95% CI)
PIU	(5, 20)	8.00	3.02	0.79	7.94	3.20	8.06	2.84	1,928,820	< 0.001	0.04 (−0.02; 0.10)
PSU	(9, 45)	23.36	8.24	0.86	22.36	8.61	24.33	7.74	1,750,946	< 0.001	0.24 (0.18; 0.30)
Externalizing	(10, 30)	16.67	3.27	0.66	16.53	3.22	16.81	3.31	1,981,714	0.018	0.09 (0.02; 0.15)
Internalizing	(10, 30)	15.38	3.35	0.63	14.63	3.13	16.12	3.40	1,523,921	< 0.001	0.45 (0.39; 0.52)
FoMO	(5, 25)	12.68	4.01	0.73	11.99	4.21	13.36	3.68	1,623,190	< 0.001	0.35 (0.29; 0.41)
Resilience	(12, 36)	29.29	3.68	0.68	29.27	3.59	29.30	3.77	2,042,408	0.434	0.01 (−0.05; 0.07)
Hopelessness	(5, 10)	5.85	1.38	0.79	5.75	1.27	5.95	1.47	1,965,174	0.001	0.14 (0.08; 0.20)

PIU, problematic internet use; PSU, problematic smartphone use; FoMO, fear of missing out; W, Wilcoxon rank sum test statistic; d, Cohen's d group differences effect size statistic

higher score indicated more severe symptoms of PSU. Of note, because the study was conducted in adolescents of whom most were not financially independent of their parents/caretakers, the last item of MPPUS-10 (“I have received mobile phone bills I could not afford to pay”) was not included in the analyses. Hence, the final variable was created as the sum of nine items.

Internalizing problems and externalizing problems were assessed using the Strengths and Difficulties Questionnaire [SDQ; 63]. The original questionnaire consisted of 25 items, but the prosocial behavior subscale was omitted from the HBSC, so the scale consisted of 20 items that covered four subscales: emotional symptoms (i.e., “I am often unhappy, downhearted or tearful”), conduct problems (i.e., “I get very angry and often lose my temper”), hyperactivity (i.e., “I am restless; I cannot stay still for long”), and peer relationship problems (i.e., “Other children or young people pick on me or bully me”). Five items were reverse coded and rescaled for further analysis. Each item had three answer options: 1 = “not true”, 2 = “somewhat true”, and 3 = “certainly true”, with a higher score indicating more internalizing of problems or externalizing of problems. Instead of four individual subscales we decided to work with broader internalizing and externalizing problem subscales because they were shown to work better in generalized (nonclinical) populations [64]. Therefore, we combined emotional and peer subscales into an internalizing problem subscale, and behavioral and hyperactivity subscales into an externalizing problem subscale. Both variables were calculated as the sum of the scores for each item.

Fear of missing out was measured using a shortened version of the Fear of Missing Out Scale [FoMO; 46]. Unlike the original 10-item questionnaire, we used a five-item version for each statement, ranging from 1 = “strongly disagree” to 5 = “strongly agree.” The final variable was computed as the sum of the scores for the five items.

Hopelessness was measured using the Hopelessness Scale for Children [HSC; 65], which is a five-item tool with answer categories 1 = “agree” and 2 = “disagree” (e.g., “All I see ahead of me are bad things, not good things”). The sum of the scores for the five items was computed.

Resilience was measured using the shortened version of the Child and Youth Resilience Measure [CYRM-12; 66]. This measure is based on the socio-ecological definition of resilience, which implies that individual, peer, family, school, and community resources contribute to positive outcomes for youth. The scale consists of 12 items that cover all previous factors (e.g., “Do you have chances to show others that you are growing up and can do things by yourself?”). For each item, participants chose between three options: 1 = “no”; 2 = “sometimes”; and 3 = “yes”. The sum of the scores of the 12 items was calculated.

Analysis

Data were analyzed using the R software v4.1.3 [67]. First, we analyzed the missing data. As mentioned previously, the participants who did not respond to at least 75% of the items on each scale were excluded from the analyses. For the rest of the sample, if there was missing data, the data were imputed using predictive mean matching with the mice package v3.14.0 [68]. Internal consistency statistics, Fisher's r-to-z transformation, which is based on correlation-difference testing (for correlations that included either PIU or PSU), and Cohen's d-s were computed using the functions in the psych package v2.2.3 [69]. Pearson's correlation coefficients were computed as statistics for the associations between variables of interest. The Wilcoxon rank-sum test was used to compute the mean differences between boys and girls.

To evaluate the complexity of the associations between psychological variables and PIU and PSU, we estimated two Gaussian graphical models [GGM; 70] with the summed scores for PSU and PIU, and other psychological variables. The plot of networks includes edges and

nodes; the former depicts association strength and direction (positive or negative), whereas the latter marks the variables. According to Rodebaugh et al. [71], the strongest nodes are those that have the most relationships with other variables in the networks such that a change in those central nodes would have a significant impact on changes in all other variables. Researchers have previously suggested that the strength of a node is a crucial index for identifying variables for developing the most effective interventions [72]. The edges in GGM are conditionally dependent relationships between the nodes. The graphical least-absolute shrinkage and selection operator, in combination with the Extended Bayesian Information Criterion (EBICglasso) model selection was used to estimate GGM [73] for parsimonious/sparse networks. In addition, all nodes were predicted by other nodes for node predictability statistics. To assess the accuracy of the network centrality estimates, case-drop bootstrapping (over 1000 permutations) was computed, and bootstrapped difference tests were run to test the differences in edge weights and node centrality. Finally, we computed the centrality stability (CS) coefficients for both models; a large coefficient indicates that the estimated centrality measure is robust [74]. The packages *qgraph* v1.9.2 [75], *bootnet* v1.5 [74], and *mgm* v1.2.12 [76] were used for network analysis.

Results

Descriptive statistics and correlation analysis

Descriptive statistics and correlation analysis results are presented in Tables 1 and 2, respectively. The correlations for boys and girls are shown separately in Additional file 4: Table S1.

Differences between boys and girls in the mean values of all of the variables were tested using the Wilcoxon rank sum test. As shown in Table 1, girls scored significantly higher on all measured variables, except for resilience,

where the difference between boys and girls was not significant. The results in Table 2 show that PIU and PSU are moderately positively correlated. Furthermore, both PIU and PSU were positively correlated with both externalizing and internalizing symptoms, FoMO and hopelessness scores with small-to-moderate effect sizes. A small negative correlation was also observed between resilience and both PSU and PIU.

Differences in correlations between PIU and PSU and other variables were tested using Fisher’s r-to-z transformation. Because the correlations were tested using the dependent sample, paired correlation-difference testing was used. In the total sample, PIU showed significantly stronger positive correlations with both externalizing and internalizing symptoms and hopelessness and a stronger but negative correlation with resilience, while PSU showed a stronger correlation with FoMO. These results are consistent with the values for the separate samples of girls and boys.

Table 3 presents the results of the unpaired correlation-difference tests between boys and girls in PIU and PSU associations with other variables. In the sample of boys, the correlation between PIU and PSU was weak. In the sample of girls these two variables correlate moderately strongly, and the difference between correlation values is significant. In the case of PIU, we observed gender differences with regards to FoMO; for girls, the correlation between PIU and FoMO was significantly stronger than that for boys. In PSU, its correlation with other variables is, in all cases, significantly stronger in the sample of girls, except for FoMO.

Network analysis

To investigate the associations between PIU, PSU, and other psychological variables in a more complex framework, we computed two regularized partial correlation networks, which involved PIU and PSU, for the total

Table 2 Correlation analysis results for the total sample

Variable	Total sample (N = 4070)							
	1	2	Paired correlation difference test (PIU-PSU)		3	4	5	6
			T	p				
1. PIU	–							
2. PSU	0.298***	–						
3. Externalizing	0.344***	0.230***	6.541	<0.001	–			
4. Internalizing	0.275***	0.200***	4.208	<0.001	0.416***	–		
5. FoMO	0.206***	0.308***	–5.775	<0.001	0.247***	0.238***	–	
6. Resilience	–0.244***	–0.076***	–9.298	<0.001	–0.405***	–0.405***	–0.094***	–
7. Hopelessness	0.262***	0.149***	6.292	<0.001	0.321***	0.423***	0.176***	–0.426***

PIU, problematic internet use; PSU, problematic smartphone use; FoMO, fear of missing out

***p < 0.001

Table 3 Correlation differences between boys and girls in associations including PIU and PSU

Variable	PIU				PSU			
	Boys	Girls	z	p	Boys	Girls	z	p
PIU	–	–	–	–	0.190	0.430	8.528	<0.001
PSU	0.190	0.430	8.528	<0.001	–	–	–	–
Externalizing	0.328	0.361	1.194	0.233	0.172	0.287	3.875	<0.001
Internalizing	0.252	0.306	1.867	0.062	0.148	0.212	2.109	0.035
FoMO	0.161	0.261	3.339	0.001	0.270	0.325	1.924	0.054
Resilience	–0.250	–0.240	0.339	0.735	–0.040	–0.115	2.406	0.016
Hopelessness	0.248	0.277	0.993	0.321	0.106	0.178	2.343	0.019

PIU, problematic internet use; PSU, problematic smartphone use; FoMO, fear of missing out. Statistically significant correlation differences are highlighted in bold font

sample and for boys and girls separately, to determine whether these specific variables were differentially associated with psychological variables. These networks are depicted in Figs. 1, 2 and 3. The edge weights of the edges of the networks in Figs. 1, 2 and 3 are in Additional file 5: Table S2.

Figures 1, 2 and 3 show that, in a large part, the networks appear very similar. One distinction between boys’ and girls’ networks is that in boys, resilience has a small negative association with PIU, whereas in girls, there is a slight positive association between resilience and PSU. The average node predictability for networks, including




Fig. 1 Regularized partial correlation networks that include either PIU (a) or PSU (b) in association with other psychological variables in the total sample. Notes: Blue lines represent positive regularized partial correlations and red lines represent negative regularized partial correlations. The line thickness indicates the strength of the relationship. The grey pie chart that surrounds each node depicts the proportion of a given node’s variance, as explained by the other nodes in the network. PIU, problematic internet use; PSU, Problematic smartphone use; FoMO, fear of missing out; INT, Internalizing; EXT, Externalizing; HOPE, Hopelessness; RES, Resilience




Fig. 2 Regularized partial correlation networks that include either PIU (a) or PSU (b) in association with other psychological variables in the boys’ sample. Notes: Blue lines represent positive regularized partial correlations and red lines represent negative regularized partial correlations. The line thickness indicates the strength of the relationship. The grey pie chart surrounding each node depicts the proportion of a given node’s variance, as explained by the other nodes in the network. PIU, problematic internet use; PSU, Problematic smartphone use; FoMO, fear of missing out; INT, Internalizing; EXT, Externalizing; HOPE, Hopelessness; RES, Resilience

PIU and PSU, was $R^2=0.241$ and $R^2=0.238$, respectively. The average node predictability statistics for the boys’ and girls’ networks were roughly of similar magnitude, with $R^2_{PIU}=0.216$ and $R^2_{PSU}=0.208$ for boys, and $R^2_{PIU}=0.265$ and $R^2_{PSU}=0.262$ for girls.

In all cases, the networks showed acceptable stability, with a centrality stability coefficient of $CS \geq 0.70$. The node strengths of these models are shown in Fig. 4. Figure 4 shows the PSU and PIU cannot be characterized as central nodes in any network. Potential differences in node centralities were also observed. Specifically, in the boys’ sample (2a and 2b tabs in Fig. 4), the most central nodes were for externalizing symptoms and resilience, while, among girls, the most central nodes seemed to be both the externalizing and internalizing symptoms.

Additional network statistics (edge weights and node strength difference test results) are presented in Additional files 1, 2 and 3. In general, it could be observed that most of the node strengths were statistically significantly different from each other. However, in all models, the externalizing and internalizing factor nodes had the

highest node strength values, and these factors were not statistically different from each other. With regard to the edge difference test results, most of the edges were statistically significantly different from each other. However, in networks involving PIU, the edges PIU-hopelessness and PIU-externalizing factors were not statistically different from each other.

Discussion

The current study aimed to investigate the associations between adolescents’ PIU, PSU, and related protective and risk factors. Specifically, we examined the extent to which PIU and PSU share the same risk and protective factors, and how these variables are interrelated in boys and girls. The findings showed that PIU and PSU were positively correlated; however, this association was weak in boys and moderate in girls. PIU and PSU showed a roughly similar structure of relationships with other variables—they were both positively associated with psychological risk factors. However, the correlations between PIU and the other variables were significantly stronger




Fig. 3 Regularized partial correlation networks that include either PIU (a) or PSU (b) in association with other psychological variables in the girls’ sample. Notes: Blue lines represent positive regularized partial correlations and red lines represent negative regularized partial correlations. The line thickness indicates the strength of this relationship. The grey pie chart surrounding each node depicts the proportion of a given node’s variance, as explained by the other nodes in the network. PIU, problematic internet use; PSU, Problematic smartphone use; FoMO, fear of missing out; INT, Internalizing; EXT, Externalizing; HOPE, Hopelessness; RES, Resilience

than those between the same variables and PSU, with FoMO as the only exception. The relationships studied also differed between boys and girls.

There can be several reasons why the associations between PIU and psychological variables were stronger than those between PSU and psychological factors. One potential reason is that children may have more access to internet-based activities via devices other than smartphones (e.g., PC, tablets, etc.). This may result in more uninterrupted time spent online, as it may be plausible that smartphones prompt interruptive notifications more frequently than, say, tablets. Owing to fewer interruptions, children may have extended their screen time with an activity. It should also be noted that when it comes to assessing internet use, the line between the use of online functionalities of a smartphone may be implicitly included in the evaluation, as PIU may be an umbrella concept covering other online-based problematic behaviors [77, 78]. Another potential explanation could be that internet use may lead to a sense of anonymity when not performed on a smartphone. In other words, one could

hypothesize that a smartphone may be associated with reduced online disinhibition [79], because communication with disclosed contacts may create a feeling of lower anonymity. Online anonymity, in turn, may promote lurking behavior—socially passive internet consumption—which has been shown to be associated with reduced mental health and problematic social media use [80, 81].

Based on these results, we cannot claim that the PIU and PSU exhibit the same phenomenon. Their mutual correlation was relatively low and they shared approximately 8.8% of the variance in the total sample. In contrast, our study showed that PIU and PSU had very similar relationships with the psychological variables. They both showed a positive association with fear of missing out, hopelessness, externalizing problems, and internalizing problems, and a negative association with resilience. These variables clearly contributed to the shared variance between PIU and PSU. This is consistent with previous studies that showed that fear of missing out, externalizing problems (i.e., impulsivity,




Fig. 4 Node strengths for the networks depicted in Fig. 1, 2 and 3. PIU, problematic internet use; PSU, Problematic smartphone use; FoMO, fear of missing out; INT, Internalizing; EXT, Externalizing; HOPELESS, Hopelessness

hyperactivity, aggression), and internalizing problems (i.e., various emotional difficulties) could be risk factors for various forms of problematic online behavior [34, 36, 49], whereas resilience is a protective factor in these cases [55, 56]. However, it must be noted that the associations between selected variables and PIU were stronger than their associations with PSU, which is in line with the study by Jeong et al. [82], who found that the risk factors for PIU were different from those of PSU and non-addicted groups. These results raise questions regarding the extent to which PSU is an independent pathological phenomenon. The only variable, whose association with PSU was stronger than with PIU, was the fear of missing out. It was previously found that people who scored high on FoMO had a higher tendency to overuse their smartphones to satisfy their need for constant connectedness [83]. Owing to their portability, smartphones can provide

24/7 internet access allowing users to constantly check what is happening online. At the same time, this permanent connectedness heightens the awareness of possibly missing out on potentially more rewarding experiences, which could fuel FoMO even more [84].

Our findings also indicate interesting differences between girls and boys. With the exception of resilience, girls showed significantly higher values for all measured variables than boys. There are several possible reasons why the associations between PIU, PSU, and psychological variables are generally stronger in girls than boys. It should be noted that similar results have been demonstrated before; specifically, girls tend to spend more time online (and on digital devices) than boys, and problematic digital technology use has also been reported higher in girls than boys [85]. Given that girls place greater importance on social relationships and it also affects

their mental well-being more than in boys [86] it is also natural that, for instance, social media usage patterns differ across genders [87]. It has been demonstrated that girls are more affected by online social comparisons [88], which could affect their body image [89]. This could, subsequently, also affect other aspects of mental health.

As for PSU, however, current research states that gender differences are not as evident, although some authors have reported that females are more susceptible to PSU [90, 91]. Recently, it has been found that girls use SNS and other social communication channels much more intensively, such as Facebook Messenger and WhatsApp [92, 93]. These applications are available predominantly on smartphones, which could partially explain why girls exhibit higher PSU values than boys do. In the case of PIU, the latest research predominantly states that boys have higher levels of PIU than girls [e.g., 94, 95], while others are in line with our findings [e.g., 96–98]. For example, Ha and Hwang [96] found that girls with emotional difficulties had a higher risk of developing internet addiction than boys with the same conditions. As the girls in our sample reported higher levels of emotional problems than boys, it is possible that these factors may have been related to more PIU. Girls also scored significantly higher on all psychological variables, except resilience, for which no difference was found between boys and girls. This result is consistent with previous studies that repeatedly report a higher prevalence for mental health problems in girls compared to boys [99–101]. During adolescence, girls are more susceptible to specific stressors associated with increased psychological distress and an increased likelihood of mental health problems, such as body dissatisfaction [102], low self-esteem [99], and academic stress and worries about school performance [103]. According to some authors [104, 105] boys may have more difficulties acknowledging and describing their mental health issues and, in comparison to girls, tend to mask or downplay their problems. This may be related to cultural expectations related to gender roles—in many societies, boys are discouraged from showing vulnerability or weakness and, thus, tend to complain less often about their health problems in general [106].

The results of the network analysis also showed that PIU and PSU were not the central nodes in any network. For boys, the node with the highest strength was externalizing problems, whereas for girls, resilience, externalizing problems, and internalizing problems were of comparable importance. In the present study, this could mean that, if the central nodes (e.g., resilience, externalizing problems, internalizing problems) are targeted, they can significantly change the levels of PIU and PSU; thus, they are ideal targets for prevention and treatment. The importance of internalizing and externalizing problems

as risk factors for the development of PIU or PSU has been demonstrated previously. For example, internalizing problems, depression, anxiety, and peer-relationship difficulties predicted both PIU and PSU in previous studies [36, 37, 56, 107]. In the case of externalizing problems, impulsivity, aggression, and attention deficit hyperactivity symptoms have been previously identified as risk factors for PIU or PSU [56, 107–110]. Some studies [101, 111] also suggest that boys have a higher tendency to externalize problems and girls internalize problems, which could explain why internalizing problems are the central node in girls' networks but not in boys' networks. At the same time, resilience has been reported to be one of the most important protective factors against the development of PIU or PSU [54, 55].

The results of our study can be used to develop intervention programs to prevent PIU and PSU. As mentioned above, targeting central nodes in a network may lead to improved mental health. As an example, externalizing symptoms were among the nodes with the highest strength in both boys and girls, meaning indicating that targeting these symptoms may lead to improved well-being in children. There are several examples of how externalizing symptoms can be addressed in children. For instance, parent training programs could be useful in teaching how to cope with children's behavior by communicating clear expectations, providing consistent consequences for misbehavior, and reinforcing positive behaviors [112]. Social skills training may also have beneficial effects on externalizing symptoms [113]. Finally, school-level interventions (e.g., mental health literacy and stigma mitigation) could also be useful in reducing the severity of externalizing symptoms [114].

This study has several limitations. First, because the design was cross-sectional, causal relationships between variables could not be inferred based on the results. Second, the study design only entailed an interindividual perspective. Previous research on individual differences, however, demonstrated that the structure and associations of inter- and intraindividual differences might not necessarily be the same [e.g., 115]; thus, future studies should incorporate an intraindividual perspective next to the interindividual perspective [116]. Third, this study did not include other factors, such as family, in the present analyses. It has previously been demonstrated that children's family circumstances, such as parental education [117], might be associated with children's digital device use [118]. Future research should consider intraindividual differences in PIU and PSU within this context. Fourth, the study used self-reported data, which might be prone to response bias such as social desirability or acquiescence. Fifth, since the data came from a complex epidemiological study, it was not always possible to use

full-length questionnaires; therefore, shortened versions of most scales were used instead. This could have negatively affected the reliability of the scales. At the same time, both of the PIU and PSU scales are designed to measure generalized internet and smartphone addiction; therefore, they cannot provide information about specific types of internet or smartphone usage behaviors (e.g., social networking, gaming, and online shopping). Despite these limitations, the key advantage of our study was its large and nationally representative sample of adolescents.

Conclusion

PIU and PSU are weakly to moderately related phenomena, yet they are distinct constructs that differ at the psychological level, with psychological risk factors mostly being especially relevant for PIU. Moreover, these phenomena were rather different between boys and girls, with stronger associations between PIU and PSU and psychological risk factors in girls.

Abbreviations

CS	Centrality stability
CYRM-12	Child and Youth Resilience Measure
EIU	Excessive Internet Use Scale
EBICglasso	Extended Bayesian Information Criterion
FoMO	Fear of missing out
GGM	Gaussian graphical models
HBSC	WHO-collaborative Health Behavior in School-aged Children study
HSC	Hopelessness Scale for Children
MPPUS-10	Problematic Mobile Phone Use Scale
PIU	Problematic internet use
PSU	Problematic smartphone use
SDQ	Strengths and Difficulties Questionnaire
SNS	Social networking sites

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13034-023-00620-z>.

Additional file 1. Supplementary Figure S1: Bootstrapped edge weights (95% CIs)

Additional file 2. Supplementary Figure S2: Edge weights differences test results

Additional file 3. Supplementary Figure S3: Centrality difference test results

Additional file 4. Supplementary Table S1: Correlation analysis results for the total sample, boys, and girls

Additional file 5. Supplementary Table S2: Edges

Acknowledgements

Not applicable.

Author contributions

DR contributed to the study conceptualization and design, planned and carried out the data analysis, interpreted the data, and drafted and revised the manuscript. LB acquired funding for the study, conceptualized and designed the study as well as drafted and revised the manuscript. NL reviewed and revised the manuscript. AF contributed to the first draft of the manuscript.

DH acquired funding and collected the data. CM reviewed and revised the manuscript. LB and NL serve as corresponding authors. All authors agree to be accountable for all aspects of this study. All authors have read and approved the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL. This study was supported by Masaryk University (Project MUNI/A/1525/2021) and the Slovak Research and Development Support Agency under Contract No. APVV-18-0070.

Availability of data and materials

The dataset used and analyzed during the current study are available from the corresponding author Lukas Blinka on scholarly request.

Declarations

Ethics approval and consent to participate

All procedures were performed in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and the Helsinki Declaration of 1975, as revised in 2000 (5). Informed consent was obtained from all patients for inclusion in the study. This study was approved by the Ethics Committee of the Medical Faculty at Pavol Jozef Safarik University in Kosice, Slovak Republic (16N/2017).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 22 March 2023 Accepted: 23 May 2023

Published online: 12 June 2023

References

- Livingstone S, Haddon L, Görzig A, Ólafsson K. Risks and safety on the internet: the perspective of European children: full findings and policy implications from the EU Kids Online survey of 9–16 year olds and their parents in 25 countries. LSE, EU Kids Online Network, London, UK; 2011.
- Hollis C, Livingstone S, Sonuga-Barke E. Editorial: the role of digital technology in children and young people's mental health—A triple-edged sword? *J Child Psychol Psychiatry*. 2020;61:837–41.
- Ferreira C, Ferreira H, Vieira MJ, Costeira M, Branco L, Dias Â, Macedo L. Epidemiologia do Uso de Internet numa População Adolescente e Sua Relação com Hábitos de Sono. *Acta Med Port*. 2017;30:524–33.
- Aghasi M, Matinfar A, Golzarand M, Salari-Moghaddam A, Ebrahimpour-Koujan S. Internet use in relation to overweight and obesity: a systematic review and meta-analysis of cross-sectional studies. *Adv Nutr*. 2020;11:349–56.
- Rozgonjuk D, Täht K. To what extent does internet use affect academic performance? Using evidence from the large-scale PISA study. *Annu Rev CyberTherapy Telemed*. 2017;15:39–44.
- Jorgenson AG, Hsiao RC-J, Yen C-F. Internet addiction and other behavioral addictions. *Child Adolesc Psychiatr Clin N Am*. 2016;25:509–20.
- Kuss D, Griffiths M, Karila L, Billieux J. Internet addiction: a systematic review of epidemiological research for the last decade. *Curr Pharm Des*. 2014;20:4026–52.
- Bickham DS. Current research and viewpoints on internet addiction in adolescents. *Curr Pediatr Rep*. 2021;9:1–10.
- Sahu M, Gandhi S, Sharma MK. Mobile phone addiction among children and adolescents: a systematic review. *J Addict Nurs*. 2019;30:261–8.
- Tateno M, Teo AR, Ukai W, Kanazawa J, Katsuki R, Kubo H, Kato TA. Internet addiction, smartphone addiction, and Hikikomori trait in Japanese young adult: social isolation and social network. *Front Psychiatry*. 2019;10:455.

11. Billieux J. Problematic use of the mobile phone: a literature review and a pathways model. *Curr Psychiatry Rev.* 2012;8:299–307.
12. Fineberg N, Demetrovics Z, Stein D, et al. Manifesto for a European research network into problematic usage of the internet. *Eur Neuropsychopharmacol.* 2018;28:1232–46.
13. Heather N. Overview of addiction as a disorder of choice and future prospects. In: Heather N, Segal G, editors. *Addiction and choice.* Oxford: Oxford University Press; 2016. p. 463–82.
14. Choi S-W, Kim D-J, Choi J-S, Ahn H, Choi E-J, Song W-Y, Kim S, Youn H. Comparison of risk and protective factors associated with smartphone addiction and Internet addiction. *J Behav Addict.* 2015;4:308–14.
15. Kwon M, Lee J-Y, Won W-Y, Park J-W, Min J-A, Hahn C, Gu X, Choi J-H, Kim D-J. Development and validation of a Smartphone Addiction Scale (SAS). *PLoS ONE.* 2013;8:e56936.
16. Lachmann B, Sindermann C, Sariyska RY, Luo R, Melchers MC, Becker B, Cooper AJ, Montag C. The role of empathy and life satisfaction in internet and smartphone use disorder. *Front Psychol.* 2018;9:398.
17. Škařupová K, Ólafsson K, Blinka L. The effect of smartphone use on trends in European adolescents' excessive internet use. *Behav Inf Technol.* 2016;35:68–74.
18. Marino C, Canale N, Melodia F, Spada MM, Vieno A. The overlap between problematic smartphone use and problematic social media use: a systematic review. *Curr Addict Rep.* 2021;8:469–80.
19. Blinka L, Škařupová K, Ševčíková A, Wölfling K, Müller KW, Dreier M. Excessive internet use in European adolescents: What determines differences in severity? *Int J Public Health.* 2015;60:249–56.
20. Rozgonjuk D, Davis KL, Montag C. The roles of primary emotional systems and need satisfaction in problematic Internet and smartphone use: a network perspective. *Front Psychol.* 2021;12:709805.
21. Mok JY, Choi S-W, Kim D-J, Choi J-S, Lee J-W, Ahn H-J, Choi E-J, Song W-Y. Latent class analysis on internet and smartphone addiction in college students. *Neuropsychiatr Dis Treat.* 2014;8:17:1.
22. Lachmann B, Duke É, Sariyska R, Montag C. Who's addicted to the smartphone and/or the Internet? *Psychol Pop Media Cult.* 2019;8:182–9.
23. Ho RC, Zhang MW, Tsang TY, et al. The association between internet addiction and psychiatric co-morbidity: a meta-analysis. *BMC Psychiatry.* 2014;14:183.
24. Koo HJ, Kwon J-H. Risk and protective factors of internet addiction: a meta-analysis of empirical studies in Korea. *Yonsei Med J.* 2014;55:1691.
25. Lam LT. Risk factors of internet addiction and the health effect of internet addiction on adolescents: a systematic review of longitudinal and prospective studies. *Curr Psychiatry Rep.* 2014;16:508.
26. Cho H-Y, Kim DJ, Park JW. Stress and adult smartphone addiction: mediation by self-control, neuroticism, and extraversion. *Stress Health.* 2017;33:624–30.
27. Sage M, Randolph K, Fitch D, Sage T. Internet use and resilience in adolescents: a systematic review. *Res Soc Work Pract.* 2021;31:171–9.
28. Lee RSC, Hoppenbrouwers S, Franken I. A systematic meta-review of impulsivity and compulsivity in addictive behaviors. *Neuropsychol Rev.* 2019;29:14–26.
29. Zhou Z, Zhu H, Li C, Wang J. Internet addictive individuals share impulsivity and executive dysfunction with alcohol-dependent patients. *Front Behav Neurosci.* 2014. <https://doi.org/10.3389/fnbeh.2014.00288>.
30. Modecki KL, Zimmer-Gembeck MJ, Guerra N. Emotion regulation, coping, and decision making: three linked skills for preventing externalizing problems in adolescence. *Child Dev.* 2017;88:417–26.
31. Cerutti R, Spensieri V, Presaghi F, Valastro C, Fontana A, Guidetti V. An exploratory study on internet addiction, somatic symptoms and emotional and behavioral functioning in school-aged adolescents. *Clin Neuropsychiatry.* 2017;14:374–83.
32. Milani L, La Torre G, Fiore M, Grumi S, Gentile DA, Ferrante M, Miccoli S, Di Blasio P. Internet gaming addiction in adolescence: risk factors and maladjustment correlates. *Int J Ment Health Addict.* 2018;16:888–904.
33. Ozturk FO, Ekinçi M, Ozturk O, Canan F. The relationship of affective temperament and emotional-behavioral difficulties to internet addiction in Turkish teenagers. *ISRN Psychiatry.* 2013;2013:1–6.
34. Riehm KE, Feder KA, Tormohlen KN, Crum RM, Young AS, Green KM, Pacek LR, La Flair LN, Mojtabai R. Associations between time spent using social media and internalizing and externalizing problems among US youth. *JAMA Psychiat.* 2019;76:1266.
35. Durbéej N, Sörman K, Norén Selinus E, Lundström S, Lichtenstein P, Hellner C, Halldner L. Trends in childhood and adolescent internalizing symptoms: results from Swedish population based twin cohorts. *BMC Psychol.* 2019;7:50.
36. Kim S-G, Park J, Kim H-T, Pan Z, Lee Y, McIntyre RS. The relationship between smartphone addiction and symptoms of depression, anxiety, and attention-deficit/hyperactivity in South Korean adolescents. *Ann Gen Psychiatry.* 2019;18:1.
37. Leo K, Kewitz S, Wartberg L, Lindenberg K. Depression and social anxiety predict internet use disorder symptoms in children and adolescents at 12-month follow-up: results from a longitudinal study. *Front Psychol.* 2021;12:787162.
38. Sevelko K, Bischof G, Bischof A, Besser B, John U, Meyer C, Rumpf H-J. The role of self-esteem in Internet addiction within the context of comorbid mental disorders: findings from a general population-based sample. *J Behav Addict.* 2018;7:976–84.
39. Velez-moro R, Lacefield K, Roberti JW. Perceived stress, sensation seeking, and college students' abuse of the Internet. *Comput Hum Behav.* 2010;26:1526–30.
40. Matar Boumosleh J, Jaalouk D. Depression, anxiety, and smartphone addiction in university students—A cross sectional study. *PLoS ONE.* 2017;12:e0182239.
41. Rębisz S, Sikora I. Internet addiction in adolescents. *Educ Pract Theory.* 2016;11:194–204.
42. Danneel S, Nelemans S, Spithoven A, Bastin M, Bijttebier P, Colpin H, Van Den Noortgate W, Van Leeuwen K, Verschuere K, Goossens L. Internalizing problems in adolescence: linking loneliness, social anxiety symptoms, and depressive symptoms over time. *J Abnorm Child Psychol.* 2019;47:1691–705.
43. Liu J, Bowker JC, Coplan RJ, Yang P, Li D, Chen X. Evaluating links among shyness, peer relations, and internalizing problems in Chinese young adolescents. *J Res Adolesc.* 2019;29:696–709.
44. Đurišić M, Gajić J. Social functioning of students with internalizing behavioral problems. *Res Pedagogy.* 2016;6:32–42.
45. Beyens I, Frison E, Eggermont S. "I don't want to miss a thing": adolescents' fear of missing out and its relationship to adolescents' social needs, Facebook use, and Facebook related stress. *Comput Hum Behav.* 2016;64:1–8.
46. Przybylski AK, Murayama K, DeHaan CR, Gladwell V. Motivational, emotional, and behavioral correlates of fear of missing out. *Comput Hum Behav.* 2013;29:1841–8.
47. Tomczyk Ł, Selmanagic-Lizde E. Fear of missing out (FOMO) among youth in Bosnia and Herzegovina—scale and selected mechanisms. *Child Youth Serv Rev.* 2018;88:541–9.
48. Gil F, Chamarro A, Oberst U. Addiction to online social networks: A question of "Fear of Missing Out"? *J Behav Addict.* 2016;4:51.
49. Alt D, Boniel-Nissim M. Using multidimensional scaling and PLS-SEM to assess the relationships between personality traits, problematic internet use, and fear of missing out. *Behav Inf Technol.* 2018;37:1264–76.
50. Elhai JD, Levine JC, Dvorak RD, Hall BJ. Fear of missing out, need for touch, anxiety and depression are related to problematic smartphone use. *Comput Hum Behav.* 2016;63:509–16.
51. Southwick SM, Charney DS. The science of resilience: implications for the prevention and treatment of depression. *Science.* 2012;338:79–82.
52. Mayordomo T, Viguier P, Sales A, Satorres E, Meléndez JC. Resilience and coping as predictors of well-being in adults. *J Psychol.* 2016;150:809–21.
53. Bilgin O, Taş I. Effects of perceived social support and psychological resilience on social media addiction among university students. *Univers J Educ Res.* 2018;6:751–8.
54. Liao YQ, Ye BJ, Jin P, Xu Q, Li AM. The effect of resilience on mobile phone addiction among minority preparatory students in Han District: moderated mediating effect. *Psychol Dev Educ.* 2017;33:487–95.
55. Robertson TW, Yan Z, Rapoza KA. Is resilience a protective factor of internet addiction? *Comput Hum Behav.* 2018;78:255–60.
56. Zhou P, Zhang C, Liu J, Wang Z. The relationship between resilience and internet addiction: a multiple mediation model through peer relationship and depression. *Cyberpsychol Behav Soc Netw.* 2017;20:634–9.
57. Li D, Zhang W, Li X, Zhen S, Wang Y. Stressful life events and problematic internet use by adolescent females and males: a mediated moderation model. *Comput Hum Behav.* 2010;26:1199–207.

58. Inchley J, Currie D, Cosma A, Smaidal O. Health behaviour in school-aged children (HBSC) study protocol: background, methodology and mandatory items for the 2017/18 survey. St Andrews: CAHRU; 2018.
59. Škařupová K, Olafsson K, Blińska L. Excessive internet use and its association with negative experiences: quasi-validation of a short scale in 25 European countries. *Comput Hum Behav*. 2015;53:118–23.
60. Griffiths M. A 'components' model of addiction within a biopsychosocial framework. *J Subst Use*. 2005;10:191–7.
61. Foerster M, Roser K, Schoeni A, Rössli M. Problematic mobile phone use in adolescents: derivation of a short scale MPPUS-10. *Int J Public Health*. 2015;60:277–86.
62. López-Fernández O, Honrubia-Serrano ML, Freixa-Blanxart M. Adaptación española del "Mobile Phone Problem Use Scale" para población adolescente. *Adicciones*. 2012;24:123.
63. Goodman R. The strengths and difficulties questionnaire: a research note. *J Child Psychol Psychiatry*. 1997;38:581–6.
64. Goodman A, Lamping DL, Ploubidis GB. When to use broader internalising and externalising subscales instead of the hypothesised five subscales on the Strengths and Difficulties Questionnaire (SDQ): data from British parents, teachers and children. *J Abnorm Child Psychol*. 2010;38:1179–91.
65. Kazdin AE, French NH, Unis AS, Esveldt-Dawson K, Sherick RB. Hopelessness, depression, and suicidal intent among psychiatrically disturbed inpatient children. *J Consult Clin Psychol*. 1983;51:504–10.
66. Liebenberg L, Ungar M, LeBlanc JC. The CYRM-12: a brief measure of resilience. *Can J Public Health*. 2013;104:e131–5.
67. R Core Team. R: a language and environment for statistical computing; 2021.
68. Van Buuren S, Groothuis-Oudshoorn K. mice: multivariate imputation by chained equations in R. *J Stat Soft*. 2011. <https://doi.org/10.18637/jss.v045.i03>.
69. Revelle W. Psych: procedures for personality and psychological research. Evanston: Northwestern University; 2021.
70. Epskamp S, Waldorp LJ, Möttus R, Borsboom D. The Gaussian graphical model in cross-sectional and time-series data. *Multivariate Behav Res*. 2018;53:453–80.
71. Rodebaugh TL, Tonge NA, Piccirillo ML, et al. Does centrality in a cross-sectional network suggest intervention targets for social anxiety disorder? *J Consult and Clin Psychol*. 2018;86:831–44.
72. Costantini G, Epskamp S, Borsboom D, Perugini M, Möttus R, Waldorp LJ, Cramer AOJ. State of the aRT personality research: a tutorial on network analysis of personality data in R. *J Res Pers*. 2015;54:13–29.
73. Epskamp S, Fried EI. A tutorial on regularized partial correlation networks. *Psychol Methods*. 2018;23:617–34.
74. Epskamp S, Borsboom D, Fried EI. Estimating psychological networks and their accuracy: a tutorial paper. *Behav Res*. 2018;50:195–212.
75. Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. Qgraph: network visualizations of relationships in psychometric data. *J Stat Soft*. 2012. <https://doi.org/10.18637/jss.v048.i04>.
76. Haslbeck JMB, Waldorp LJ. Mgm: estimating time-varying mixed graphical models in high-dimensional data; 2015. <https://doi.org/10.48550/ARXIV.1510.06871>.
77. Baggio S, Starcevic V, Studer J, Simon O, Gainsbury SM, Gmel G, Billieux J. Technology-mediated addictive behaviors constitute a spectrum of related yet distinct conditions: a network perspective. *Psychol of Addict Behav*. 2018;32:564–72.
78. Rozgonjuk D, Schivinski B, Pontes HM, Montag C. Problematic online behaviors among gamers: the links between problematic gaming, gambling, shopping, pornography use, and social networking. *Int J Ment Health Addic*. 2023;21:240–57.
79. Scott RA, Stuart J, Barber BL. What predicts online disinhibition? Examining perceptions of protection and control online and the moderating role of social anxiety. *Cyberpsychol Behav Soc Netw*. 2022;25:294–300.
80. Verduyn P, Gugushvili N, Kross E. Do social networking sites influence well-being? The extended active-passive model. *Curr Dir Psychol Sci*. 2022;31:62–8.
81. Rozgonjuk D, Ryan T, Kuljus J-K, Täht K, Scott GG. Social comparison orientation mediates the relationship between neuroticism and passive Facebook use. *Cyberpsychology*. 2019. <https://doi.org/10.5817/CP2019-1-2>.
82. Jeong B, Lee JY, Kim BM, Park E, Kwon J-G, Kim D-J, Lee Y, Choi J-S, Lee D. Associations of personality and clinical characteristics with excessive Internet and smartphone use in adolescents: a structural equation modeling approach. *Addict Behav*. 2020;110:106485.
83. Rozgonjuk D, Sindermann C, Elhai JD, Montag C. Fear of missing out (FoMO) and social media's impact on daily-life and productivity at work: Do WhatsApp, Facebook, Instagram, and Snapchat use disorders mediate that association? *Addict Behav*. 2020;110:106487.
84. Milyavskaya M, Saffran M, Hope N, Koestner R. Fear of missing out: prevalence, dynamics, and consequences of experiencing FoMO. *Motiv Emot*. 2018;42:725–37.
85. Twenge JM, Martin GN. Gender differences in associations between digital media use and psychological well-being: Evidence from three large datasets. *J Adolesc*. 2020;79:91–102.
86. Flook L. Gender differences in adolescents' daily interpersonal events and well-being. *Child Dev*. 2011;82:454–61.
87. Yau JC, Reich SM. "It's just a lot of work": adolescents' self-presentation norms and practices on Facebook and Instagram. *J Res Adolesc*. 2019;29:196–209.
88. Fox J, Vendemia MA. Selective self-presentation and social comparison through photographs on social networking sites. *Cyberpsychol Behav Soc Netw*. 2016;19:593–600.
89. Hogue JV, Mills JS. The effects of active social media engagement with peers on body image in young women. *Body Image*. 2019;28:1–5.
90. Lee KE, Kim S-H, Ha T-Y, Yoo Y-M, Han J-J, Jung J-H, Jang J-Y. Dependency on smartphone use and its association with anxiety in Korea. *Public Health Rep*. 2016;131:411–9.
91. Yang Z, Asbury K, Griffiths MD. Do Chinese and British university students use smartphones differently? A cross-cultural mixed methods study. *Int J Ment Health Addic*. 2019;17:644–57.
92. Dufour M, Brunelle N, Tremblay J, Leclerc D, Cousineau M-M, Khazaal Y, Légaré A-A, Rousseau M, Berbiche D. Gender difference in internet use and internet problems among Quebec high school students. *Can J Psychiatry*. 2016;61:663–8.
93. Montag C, Błaskiewicz K, Sariyska R, Lachmann B, Andone I, Trendafilov B, Eibes M, Markowitz A. Smartphone usage in the 21st century: Who is active on WhatsApp? *BMC Res Notes*. 2015;8:331.
94. Rigelsky M, Megyesi S, Ivankova V, Al Khouri I, Sevl J. Gender differences in internet addiction among university students in the Slovak Republic. *Adiktologie*. 2021;21:35–42.
95. Shan X, Ou Y, Ding Y, Yan H, Chen J, Zhao J, Guo W. Associations between internet addiction and gender, anxiety, coping styles and acceptance in university freshmen in South China. *Front Psychiatry*. 2021;12:558080.
96. Ha Y-M, Hwang WJ. Gender differences in internet addiction associated with psychological health indicators among adolescents using a national web-based survey. *Int J Ment Health Addic*. 2014;12:660–9.
97. Hetzel-Riggin MD, Pritchard JR. Predicting problematic internet use in men and women: the contributions of psychological distress, coping style, and body esteem. *Cyberpsychol Behav Soc Netw*. 2011;14:519–25.
98. Procházka R, Suchá J, Dostál D, Dominik T, Dolejš M, Šmahaj J, Kolařík M, Glaser O, Viktorová L, Friedlová M. Internet addiction among Czech adolescents. *Psych J*. 2021;10:679–87.
99. Aanesen F, Meland E, Torp S. Gender differences in subjective health complaints in adolescence: the roles of self-esteem, stress from school-work and body dissatisfaction. *Scand J Public Health*. 2017;45:389–96.
100. Campbell OLK, Bann D, Patalay P. The gender gap in adolescent mental health: a cross-national investigation of 566,829 adolescents across 73 countries. *SSM Popul Health*. 2021;13:100742.
101. Van Droogenbroeck F, Spruyt B, Keppens G. Gender differences in mental health problems among adolescents and the role of social support: results from the Belgian health interview surveys 2008 and 2013. *BMC Psychiatry*. 2018;18:6.
102. Haugen T, Johansen BT, Ommundsen Y. The role of gender in the relationship between physical activity, appearance evaluation and psychological distress. *Child Adolesc Ment Health*. 2014;19:24–30.
103. Klinger DA, Freeman JG, Bilz L, Liiv K, Ramelow D, Sebok SS, Samdal O, Dur W, Rasmussen M. Cross-national trends in perceived school pressure by gender and age from 1994 to 2010. *Eur J Public Health*. 2015;25:51–6.

104. Patel V, Flisher AJ, Hetrick S, McGorry P. Mental health of young people: a global public-health challenge. *Lancet*. 2007;369:1302–13.
105. Rice SM, Purcell R, McGorry PD. Adolescent and young adult male mental health: transforming system failures into proactive models of engagement. *J Adolesc Health*. 2018;62:S9–17.
106. Evans J, Frank B, Oliffe JL, Gregory D. Health, Illness, Men and Masculinities (HIMM): a theoretical framework for understanding men and their health. *J Men's Health*. 2011;8:7–15.
107. Wang B, Yao N, Zhou X, Liu J, Lv Z. The association between attention deficit/hyperactivity disorder and internet addiction: a systematic review and meta-analysis. *BMC Psychiatry*. 2017;17:260.
108. Lee M, Chung SJ, Lee Y, Park S, Kwon J-G, Kim DJ, Lee D, Choi J-S. Investigation of correlated internet and smartphone addiction in adolescents: copula regression analysis. *Int J Environ Res Public Health*. 2020;17:5806.
109. Panagiotidi M, Overton P. Attention deficit hyperactivity symptoms predict problematic mobile phone use. *Curr Psychol*. 2022;41:2765–71.
110. Peterka-Bonetta J, Sindermann C, Elhai JD, Montag C. Personality associations with smartphone and internet use disorder: a comparison study including links to impulsivity and social anxiety. *Front Public Health*. 2019;7:127.
111. Lau TWI, Lim CG, Acharryya S, Lim-Ashworth N, Tan YR, Fung SSD. Gender differences in externalizing and internalizing problems in Singaporean children and adolescents with attention-deficit/hyperactivity disorder. *Child Adolesc Psychiatry Ment Health*. 2021;15:3.
112. Sukhodolsky DG, Gladstone TR, Marsh CL, Cimino KR. Behavioral interventions for irritability in children and adolescents. In: Roy AK, Brotman MA, Leibenluft E, editors. *Irritability in pediatric psychopathology*. Oxford: Oxford University Press; 2019. p. 255–74.
113. Van Der Stouwe T, Gubbels J, Castenmiller YL, Van Der Zouwen M, Asscher JJ, Hoes M, Van Der Laan PH, Stams GJJM. The effectiveness of social skills training (SST) for juvenile delinquents: a meta-analytical review. *J Exp Criminol*. 2021;17:369–96.
114. Pataño B, Miller P, Baldwin R, Taylor N, Hyder S. A systematic mapping review of interventions to improve adolescent mental health literacy, attitudes and behaviours. *Early Interv Psychiatry*. 2021;15:1470–501.
115. Löchner N, Bückle S, Olderbak S. Affect-specific empathy: experience sampling and multilevel structural equation modeling provide a within-person perspective. *Emotion*. 2022. <https://doi.org/10.1037/emo0001105>.
116. Beck ED, Jackson JJ. Idiographic traits: a return to Allportian approaches to personality. *Curr Dir Psychol Sci*. 2020;29:301–8.
117. Pons M, Bannasar-Veny M, Yañez AM. Maternal education level and excessive recreational screen time in children: a mediation analysis. *Int J Environ Res Public Health*. 2020;17:8930.
118. Vaala Sarah E. Mothers' cognitions and structural life circumstances as predictors of infants' and toddlers' television and video exposure (2012). Dissertations available from ProQuest. AAI3509498. <https://repository.upenn.edu/dissertations/AAI3509498>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

